

Article

Evaluation of Bioactive Compounds and Antimicrobial Analysis of the Essential Oil From the Leaves of *Sauropus androgynous* in Rajasthan India

Alagbe, Olujimi John^{*1,2}

1. Assistant Professor, Department of Animal Nutrition and Biochemistry, Sumitra Research Institute, Gujarat, India
2. Adjunct Lecturer, Department of Animal Biochemistry, Gandhi College of Agriculture Rajasthan India

*Correspondence: dralagbe@outlook.com

Citation: John, A. O. Evaluation of Bioactive Compounds and Antimicrobial Analysis of the Essential Oil From the Leaves of *Sauropus androgynous* in Rajasthan India. American Journal of Biology and Natural Sciences 2026, 3(1), 124-131.

Received: 08th Dec 2025

Revised: 22nd Dec 2025

Accepted: 29th Dec 2025

Published: 16th Jan 2026



Copyright: © 2026 by the authors. Submitted for open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>)

Abstract: *Sauropus androgynous* essential oil contains unique bioactive compounds with therapeutic potentials and it delivers powerful antioxidant, anti-inflammatory, antidiabetic, cytotoxic, gastro-protective, immune-protective, cardio-protective, antimicrobial, antidiarrheal, anti-helminthic amongst others. GC-MS analysis of *Sauropus androgynous* essential oil revealed the presence of 33 bioactive compounds were recorded representing 96.27 % composition dominated by Squalene (19.63 %), Caryophyllene (13.92 %), Nonane, 4-methyl (13.82 %) and n-hexadecanoic acid (10.45 %). Other bioactive compounds examined had a concentration less than 3 %. A synergy between the major and minor compounds are working together to protect, heal, restore and improve the overall vitality of the body. The result obtained on the antimicrobial activity of *Sauropus androgynous* essential oil revealed a high percentage inhibition 77.81 % , 72.00 %, 80.09 %, 73.73 % and 76.26 % against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi* respectively. The results compare well with that of the standard drug (Ciproflaxacin) (65.00 -71.00 %). It was concluded that *Sauropus androgynous* essential oil contains measurable concentration of compounds that can disrupt the cell membrane of pathogens and will also help to address the increasing cases of antimicrobial resistance.

Keywords: Bioactive, Compounds, Antimicrobials, Free Radicals, Oxidative Stress

Introduction

Sauropus androgynous, an evergreen, multipurpose herb belongs to the Phyllanthaceae plant family which is found in the tropical rain forests of West Africa, although it is also widely distributed in parts of Asia, South America and Australia (Purba and Paengkoum, 2012; Bao et al., 2020). Inside its leaves are bioactive compounds that go far beyond nutrition (Juxian and Xian, 2022; Hernandez and Alagbe, 2025a). The leaves are rich in minerals like calcium, magnesium and iron supporting bone strength and electrolyte balance (Singh et al., 2011; Singh et al., 2021), vitamins (B-complex and C) and protein between 6 – 10 % (Lin et al., 1999; Basker, 2012). For centuries, Traditional healers have used its

leaves to soothe infection, skin disorder, pyrexia, bronchitis, diarrhea and relief from acute and chronic infections because its tissues are filled with therapeutic potentials (Samad et al., 2013; John, 2024b).

The plant contains a unique profile of flavonoids and phenolic acids, nature's compounds known to neutralize the activities of free radicals, reinforces the cardiovascular system by strengthening blood vessel, acts as liver protectors, metabolic regulators, reducing inflammation and restoring enzyme balance (Alagbe, 2024; Singh et al., 2022). They also contains alkaloids, tannins, steroids and saponins which possess several pharmacological functions which includes, cytotoxic, antimicrobial, antitumor, antidiabetic, anti-cancer, antifungal, immune-modulatory, gastro-protective, derma-protective, anti-helminthic, antidiarrheal, anti-allergic, antihistaminic, antipyretic, antiseptic, antispasmodic, antiulcer and antiviral (Naemsuvan, 2013; Zhang et al., 2022). Methanolic extract from *Sauropus androgynous* leaves and stems show high radical scavenging free activity (Gonzalez et al., 2022; Purba et al., 2022), measurable compounds inhibits the activities of bacteria cells in microbial assays (Caldeira et al., 2004; Chatrou et al., 2012). Its poultices can be used for skins and wounds (Gao et al., 2014; Musa et al., 2020). Aqueous extract suppresses inflammatory cytokines, reducing airway infections and regulating sugar absorption in the body (Adewale et al., 2021; John, 2024a).

Previous studies by Rajeswari et al. (2018) have shown that *Sauropus androgynous* leaf extract contained 9,12,15-Octadecatrienoic acid at 12.90 %, Ethyl 9,12,15-Octadecatrienoate (15.93 %), Ethyl (9Z,12Z)-9,12-Octadecadienoate (11.28 %), Hexadecanoic acid (13.76 %), 1-(+)-Ascorbic acid 2,6-Dihexadecanoate (27.81 %) as the prominent bioactive compounds during GC-MS analysis. Senthamarai and Anusha (2012) also reported that GC-MS profiling in the leaves of *Sauropus Androgynus* showed that it contained 2(1H) Naphthalenone (41.17 %), Azulene (36.20 %), Pyrene, hexadecahydro (9.07 %), squalene (8.06 %) and 1, 14-Tetradecanediol (2.82 %) as major bioactive compounds.

Previous report has shown that that for every newly developed antibiotic, bacterial strains rapidly develop resistance mechanisms against it. Therefore, there is an urgent need for alternative ways to treat bacterial infections and to combat the problem of antimicrobial resistance (Carranza et al. 2015; Hernandez and Alagbe, 2025b). One of the alternatives that can be utilised for combating antimicrobial resistance is traditional herbal remedies from plants. Evaluating the bioactive compounds in *Sauropus androgynous* oil will further help to unveil their pharmacological and therapeutic properties.

Materials and Methods

Location of the experiment, plant collection and extraction of *Sauropus androgynous* oil

This study was carried out at the Microbiology Department Gandhi College of Agriculture, Rajasthan India between October to December 2025. The Institution is situated between 21°35'N 26°09'E East India. *Sauropus androgynous* fresh leaves were harvested from Rajasthan India and sent to the department of botany, Gandhi College of Agriculture for identification before it was registered under voucher number HU09/2025C/008. The identified leaves were sorted and shade dried for 13 days until a constant weight was achieved. Dried *Sauropus androgynous* leaves were pulverized using mechanical grinder. Extraction of oil was done by hydrodistillation with a Clevenger-type apparatus according to the procedures outlined by Singh et al. (2021). 250 g of the pulverized *Sauropus androgynous* was added to 1000 mL of water heated in a glass flask at 60 °C for 20 minutes, steam passes via the condenser and when cooled it was collected in a beaker. The oil collected by decantation at the end of the distillation was filtered, dried over column of anhydrous sodium sulfate, and introduced into glass bottles and stored in a refrigerator at 4 °C.

Gas chromatography – mass spectrometry analysis of *Sauropus androgynous* essential oil

GC/MS analysis of *Sauropus androgynous* essential oil was done using Claudus 5006 GC-MS Auto Sampler (China) equipped with two silica capillary columns, interfaced with a quadrupole detector (single quadrupole acquisition Method-MS parameters report), source temperature 230°C, Quadrupole temperature 150 °C; the temperature program was 60 °C for 2 min, 60-240 °C at 3 °C/min, then kept at 240 °C during 8 min; injector temperature, 240 °C. The mass spectrometry transfer line temperature, 250 °C; carrier gas, helium at a flow rate of 0.7 ml/min; injection type, split, 20:1; ionization voltage, 70 eV; electron multiplier 1000 eV; scan range 33-400 amu; scan rate, 1.56 scan/s.

Identification of components Interpretation on mass spectrum GC-MS was conducted using the database of National Institute Standard and Technology (NIST, 2001) having more than 62,000 patterns. The spectrum of the unknown component was compared with the spectrum of the known components stored in the NIST library.

Antimicrobial Assay

The antimicrobial activity of essential oil of *Sauropus androgynous* essential oil was screened against five standard strains of stock bacteria from the Microbiology department, Gandhi College of Agriculture, Rajasthan. *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi*. Microplate Alamar Blue Assay was used to determine susceptibility or resistance of the essential oils to all the selected bacteria strains. Organisms were grown in Mueller Hinton broth and inoculums were adjusted to 0.5 McFarland standard. Essential oil (20 µg/mL) was added in the wells, control wells do not contain essential oil. The volume of 96-well plate was made up to 200 µL. Finally 5 x10⁶ cells were added in all wells including both control and test. The plate was sealed with parafilm and incubated for 18 - 20 hours. Alamar Blue Dye was dispensed in each well and shaken at about 80 revolution per minute in a shaking incubator for 2 – 3 hours. Plates were covered with foil in shaking incubator. Change in color of Alamar Blue dye from blue to pink indicated the growth in bacterial strains. Ciproflaxacin was used as the reference drug.

Bioactive compounds in *Sauropus androgynous* essential oil by GC-MS analysis is presented in Table 1. A total of 33 bioactive compounds were recorded representing 96.27 % composition dominated by Squalene (19.63 %), Caryophyllene (13.92 %), Nonane, 4-methyl (13.82 %) and n-hexadecanoic acid (10.45 %). These compounds have been previously associated with anti-inflammatory (Sinha and Munshi, 2011; Namkeleja et al., 2014), antioxidant (Omokore and Alagbe, 2019; Oluwafemi et al., 2021), cytotoxic and immuno-stimulatory properties (Alagbe, 2022; Muritala et al., 2022). According to Shittu and Alagbe (2020), Squalene has the capability to neutralize free radicals activities in the body, support blood vessels integrity and protects the heart, lungs and nervous systems. Caryophyllene have been reported to be a proven powerhouse, kills pathogens and inhibits fungal growth (Tiwari et al., 2011; Ojediran et al., 2024). Hexadecanoic acid excels in lowering blood sugar and supports liver detoxification (Sharma, 2012; Jing-Chung et al., 2007; Lima et al., 2010). Nonane, 4-methyl can interfere with mitochondrion energy production in cancer cells, fights infection and calm the nervous system (Choi et al., 2013). All the 33 bioactive compounds have numerous pharmacological properties making them useful in traditional medicine for the treatment of acute and chronic inflammation, cough, fever, gastrointestinal disorders, skin infection, eye problems, sexually transmitted infections, stomach ulcers, pain, snake bite, tooth ache amongst others (Alagbe et al., 2022; Agubosi et al., 2022). A previous study by Senthamarai and Anusha (2012) on the GC-MS analysis of *Sauropus androgynous* leaf reveals that squalene and phytol contained a concentration of 8.06 % and 0.88 % respectively. Lee et al. (2011) also recorded a lower concentration of 0.12 % and 3.07 % for phytol and hexadecanoic acid. Rajeswari et al. (2018); Basker (2012) reported that *Sauropus androgynous* leaf extract contained a higher concentration of 13.80 % hexadecanoic acid. Samad et al. (2018) reported 10.40 % and 0.04 % for hexadecanoic acid and 9, 17-Octadecadienal, and these values were almost similar to the outcome in this experiment. However, geographical location, age of plant, specie, storage method, harvesting technique, processing methods have been identified as factors that influence the concentration of bioactive compounds in medicinal plants (Awa et al., 2012). A synergy between the minor and major components in essential oil from *Sauropus androgynous* in this study forms a rich network of phytochemicals that reduces inflammation by suppressing pro-inflammatory cytokines (Momza et al., 2012; Alagbe et al., 2022), produces bronchodilator effect by opening respiratory airway thereby making breathing easier (Zubairu et al., 2025), improve cardio-vascular health and electrolyte balance thus improving the overall health and vitality of the body (Alagbe, 2025).

The result obtained on the antimicrobial activity of *Sauropus androgynous* essential oil revealed a high percentage inhibition 77.81 % , 72.00 % , 80.09 % , 73.73 % and 76.26 % against *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi* respectively. The results compare well with that of the standard drug (Ciproflaxacin) (65.00 -71.00 %). The presence of squalene, the prominent bioactive compounds in the oil might have been responsible for their antimicrobial

activity. Squalene contains measurable compounds that inhibits the activities of some bacteria's (Ajiboye et al., 2013). Phyto-components like hexadecanoic acid, caryophyllene, cyclopentaneundecanoic acid, 8-octadecenoic acid, humulene, gamma.-Terpinene-3-Carene, 9, 17-Octadecadienal, Undecane, 2,10-dimethyl, 2-methyl-Glutaric acid, 2-Ethylhexyl mercaptoacetate and Linoleic acid ethyl ester have been confirmed to show antibacterial activities (Ajayi et al., 2011). This suggests that dietary supplementation of *Sauropus androgynous* essential oil can help to reduce the increasing cases of antimicrobial resistance and promote food safety (Ajayi et al., 2011). Paul et al. (2011), essential oil from *Sauropus androgynous* can defend against infection, disrupts cell membrane of pathogens and inhibited *Staphylococcus spp* and *Salmonella spp* in antimicrobial assay. Preveen et al. (2011) reported that *Sauropus androgynous* oil is potent in controlling bacterial growth in biofilms of *Streptococcus spp* and *Shigella spp*.

Table 1. Bioactive compounds in *Sauropus androgynous* essential oil by GC-MS

Compounds	R.T (min)	Molecular formula	Molecular weight (g/mol)	% Area
n-hexadecanoic acid	16.73	C ₁₆ H ₃₂ O ₂	256.424	10.45
Dipyrimadole	18.52	C ₂₄ H ₄₀ N ₈ O ₄	504.626	2.31
Octadec-9-enoic acid	18.90	C ₁₈ H ₃₄ O ₂	282.47	0.94
N-3-methylButyl acetamide	18.92	C ₇ H ₁₅ NO	129.20	1.46
Benzene,1,4-dichloro	21.13	C ₆ H ₄ Cl ₂	147.00	0.01
8-octadecenoic acid	22.46	C ₁₉ H ₃₆ O ₂	296.487	0.56
Humulene	22.78	C ₁₅ H ₂₄	204.35	0.18
Undecane, 2,10-dimethyl	22.98	C ₁₃ H ₂₈	184.36	0.46
Oxalic acid	24.06	C ₁₅ H ₂₈ O ₄	272.38	2.77
Nonane, 4-methyl	25.67	C ₁₀ H ₂₂	142.28	13.82
2-Ethylhexyl mercaptoacetate	25.75	C ₁₀ H ₂₀ O ₂ S	247.34	1.81
gamma.-Terpinene-3-Carene	26.31	C ₁₀ H ₁₆	136.23	2.28
Hexatriacontane	28.94	C ₃₆ H ₇₄	507.00	2.15
3-Isothiazolone	29.07	C ₃ H ₃ NOS	101.13	1.26
Nerolidol	29.35	C ₁₅ H ₂₆ O	222.37	0.14
10-Methylnonadecane	29.94	C ₂₀ H ₄₂	282.5	1.33
Cyclopropane	30.05	C ₃ H ₆	42.08	2.09
Ethyl palmitate	31.24	C ₁₈ H ₃₆ O ₂	284.5	0.71
2-methyl-Glutaric acid	31.88	C ₆ H ₁₀ O ₄	146.14	0.20
Linoleic acid ethyl ester	31.90	C ₁₈ H ₃₂ O ₂	308.5	0.38
9,17-Octadecadienal	33.25	C ₁₈ H ₃₂ O	264.4	0.05
4-Trifluoroacetoxytetradecane	33.57	C ₁₆ H ₂₉ F ₃ O ₂	424.4	0.57
Hexacosanoic acid	33.89	C ₂₆ H ₅₂ O ₂	396.7	1.10
Beta.-Famesene	34.21	C ₁₅ H ₂₄	204.35	1.59
Cyclopentaneundecanoic acid	34.57	C ₁₆ H ₃₀ O ₂	254.41	1.87
Pentadecane	37.21	C ₁₅ H ₃₂	212.41	0.01
Caryophyllene	37.67	C ₁₅ H ₂₄	204.351	13.92
Trans-Nerolidol	38.58	C ₁₅ H ₂₆ O	222.4	1.12
Methyl stearate	38.71	C ₁₉ H ₃₈ O ₂	298.5	1.07
Ethyl ricinoleate	39.12	C ₂₀ H ₃₈ O ₃	326.5	0.08

Squalene	39.65	C ₃₀ H ₅₀	410.7	19.63
Hexahydrofarnesyl acetone	40.19	C ₁₈ H ₃₆ O	268.5	1.24
Phytol	41.25	C ₂₂ H ₄₂ O ₂	296.5	8.71
Total				96.27
Number of compounds				33.00

Table 2. Antimicrobial activity of *Sauropus androgynous* essential oil

Microorganisms	Ciproflaxacin	% Inhibition
<i>Staphylococcus aureus</i>	71	77.81
<i>Escherichia coli</i>	65	72.00
<i>Pseudomonas aeruginosa</i>	75	80.09
<i>Shigella flexineri</i>	68	73.73
<i>Salmonella typhi</i>	70	76.26

Conclusion

It was concluded that *Sauropus androgynous* essential oil contains several unique bioactive compounds with therapeutic potentials and has been traditionally utilized for the treatment of cold, fever, inflammation, acute and chronic infections. *In vitro* studies have also shown that these compounds also have antimicrobial properties and was able to inhibit the activities of *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Shigella flexineri* and *Salmonella typhi*. Together they form a weapon against infection, inflammation and degeneration.

REFERENCES

- [1] A. P. A. Samad, U. Santoso, C. S. Chang, and F. H. Nan, "Identifying and analyzing bioactive components of *Sauropus androgynus* ethanolic leaves extract," *J. Fish. Soc. Taiwan*, vol. 40, pp. 257–265, 2013.
- [2] I. A. Ajayi, O. Ajibade, and R. A. Oderinde, "Preliminary phytochemical analysis of some plant seeds," *Res. J. Chem. Sci.*, vol. 1, no. 3, pp. 58–63, 2011.
- [3] A. A. Ajiboye, D. A. Agboola, O. Y. Fadimu, and A. O. Afolabi, "Antibacterial, phytochemical and proximate analysis of *Prosopis africana* (Linn.) seed and pod extracts," *FUTA J. Res. Sci.*, vol. 9, no. 1, pp. 101–109, 2013.
- [4] C. Q. Lin, W. B. Lin, W. D. Pan, and Y. J. Lee, "Study on chemical constituents of the essential oil from the leaves of *Sauropus androgynus* L. Merr.," *J. Trop. Subtrop. Bot.*, vol. 3, pp. 255–256, 1999.
- [5] A. Basker, "Phytochemical analysis and GC–MS profiling in the leaves of *Sauropus androgynus* (L.) Merr.," *Int. J. Drug Dev. Res.*, vol. 4, pp. 162–167, 2012.
- [6] J. Choi et al., "The antibacterial activity of various saturated and unsaturated fatty acids against several oral pathogens," *J. Environ. Biol.*, vol. 34, pp. 673–676, 2013.
- [7] U. T. Mamza, O. A. Sodipo, and I. Z. Khan, "Gas chromatography–mass spectrometry (GC–MS) analysis of bioactive components of *Phyllanthus amarus* leaves," *Int. Res. J. Plant Sci.*, vol. 3, no. 10, pp. 208–215, 2012.
- [8] E. P. Awa, S. Ibrahim, and D. A. Ameh, "GC/MS analysis and antimicrobial activity of dimethyl ether fraction of methanolic extract from stem bark of *Annona senegalensis* Pers.," *Int. J. Pharm. Sci. Res.*, vol. 3, no. 11, pp. 4213–4218, 2012.
- [9] R. Bose, M. S. Kumar, A. Manivel, and S. C. Mohan, "Chemical constituents of *Sauropus androgynus* and evaluation of its antioxidant activity," *Res. J. Phytochem.*, vol. 12, pp. 7–13, 2018.
- [10] S. Singh et al., "Estimation of proximate composition, micronutrients and phytochemical compounds in traditional vegetables from Andaman and Nicobar Islands," *Int. J. Food Sci.*, vol. 62, pp. 765–773, 2011.

- [11] J. Gu and Y. Xian, "Studies on the toxicology of *Sauropus androgynus*, a wild vegetable in South China," *J. South China Agric. Univ.*, vol. 28, pp. 10–14, 2005. [Online]. Available: <https://agris.fao.org/agris-search/search.do?recordID=CN2006002063>. Accessed: Nov. 5, 2022.
- [12] R. A. P. Purba and P. Paengkoum, "Exploring the phytochemical profiles and antioxidant, antidiabetic, and antihemolytic properties of *Sauropus androgynus* dried leaf extracts for ruminant health and production," *Molecules*, vol. 27, p. 8580, 2022, doi: 10.3390/molecules27238580.
- [13] N. Andarwulan et al., "Polyphenols, carotenoids, and ascorbic acid in underutilized medicinal vegetables," *J. Funct. Foods*, vol. 4, pp. 339–347, 2012.
- [14] H. Bao et al., "Depolymerization of lignin into monophenolics by ferrous/persulfate reagent under mild conditions," *ChemSusChem*, vol. 13, pp. 6582–6593, 2020.
- [15] R. A. P. Purba, S. Paengkoum, and P. Paengkoum, "Development of a simple high-performance liquid chromatography-based method to quantify synergistic compounds and their composition in dried leaf extracts of *Piper sarmentosum* Roxb.," *Separations*, vol. 8, p. 152, 2021.
- [16] O. Neamsuvan and T. Ruangrit, "A survey of herbal weeds that are used to treat gastrointestinal disorders from southern Thailand: Krabi and Songkhla provinces," *J. Ethnopharmacol.*, vol. 196, pp. 84–93, 2017.
- [17] A. J. A. Petrus, "*Sauropus androgynus* (L.) Merrill—a potentially nutritive functional leafy vegetable," *Asian J. Chem.*, vol. 25, pp. 9425–9433, 2013.
- [18] D. Zhang et al., "Recent trends in the micro-encapsulation of plant-derived compounds and their specific application in meat as antioxidants and antimicrobials," *Meat Sci.*, vol. 191, p. 108842, 2022.
- [19] J. González-Romero, E. J. Guerra-Hernández, and C. Rodríguez-Pérez, "Bioactive compounds from *Moringa oleifera* as promising protectors of in vivo inflammation and oxidative stress processes," in *Current Advances for Development of Functional Foods Modulating Inflammation and Oxidative Stress*, B. Hernández-Ledesma and C. Martínez-Villaluenga, Eds. Cambridge, MA, USA: Academic Press, 2022, pp. 379–399.
- [20] R. A. P. Purba, S. Paengkoum, C. Yuangklang, and P. Paengkoum, "Flavonoids and their aromatic derivatives in *Piper betle* powder promote in vitro methane mitigation in a variety of diets," *Cienc. Agrotec.*, vol. 44, p. e012420, 2020.
- [21] P. Mariya and B. A. K., "Antibacterial activity of *Sauropus androgynus* (L.) Merr.," *Int. J. Plant Sci.*, vol. 6, no. 1, pp. 189–192, 2011.
- [22] P. K. Praveen, S. Kumaravel, and C. Lalitha, "Screening of antioxidant activity, total phenolics and GC–MS study of *Vitex negundo*," *Afr. J. Biochem. Res.*, vol. 4, no. 7, pp. 191–195, 2011.
- [23] M. G. Carranza, M. B. Sevigny, D. Banerjee, and L. Fox-Cubley, "Antibacterial activity of native California medicinal plant extracts isolated from *Rhamnus californica* and *Umbellularia californica*," *Ann. Clin. Microbiol. Antimicrob.*, vol. 14, no. 1, p. 29, 2015.
- [24] I. Caldeira et al., "Improved method for extraction of aroma compounds in aged brandies and aqueous alcoholic wood extracts using ultrasound," *Anal. Chim. Acta*, vol. 513, no. 1, pp. 125–134, 2004.
- [25] L. W. Chatrou et al., "A new subfamilial and tribal classification of the pantropical flowering plant family Annonaceae informed by molecular phylogenetics," *Bot. J. Linn. Soc.*, vol. 169, no. 1, pp. 5–40, 2012.
- [26] S. Gao, G. D. Lewis, M. Ashokkumar, and Y. Hemar, "Inactivation of microorganisms by low-frequency high-power ultrasound: 1. Effect of growth phase and capsule properties of the bacteria," *Ultrason. Sonochem.*, vol. 21, no. 1, pp. 446–453, 2014.
- [27] A. O. John, "Clerodendron splendens leaf extract supplementation in weaner rabbits: Impact on growth performance, haematology and intestinal microbial population," *Cerrado: Agric. Biol. Res.*, vol. 1, no. 1, pp. 21–31, 2024.
- [28] A. O. John, "Effect of coconut shell extract on the growth performance and some haemato-biochemical parameters of broiler chicken," *Brazilian J. Sci.*, vol. 3, no. 6, pp. 82–95, 2024.

- [29] A. O. John, "Impact of dietary supplementation of *Rhamnus prinoides* leaf extract on the growth performance, nutrient retention and intestinal microbial count of Japanese quails," *Brazilian J. Sci.*, vol. 3, no. 5, pp. 40–50, 2024.
- [30] A. O. John, "Effect of performance, serum biochemistry and haematological components of feeding Japanese quails phytogenic feed additions comprising *Megaphrynium macrostachyum* leaves," *Brazilian J. Sci.*, vol. 3, no. 5, pp. 51–64, 2024.
- [31] S. Singh, O. J. Alagbe, X. Liu, R. Sharma, and A. Kumar, "Comparative analysis of ethanolic *Juniperus thurifera* leaf, stem bark and root extract using gas chromatography–mass spectrometry," *Int. J. Agric. Anim. Prod.*, vol. 2, no. 6, pp. 18–27, 2022.
- [32] A. S. Singh, J. O. Alagbe, S. Sharma, R. A. Oluwafemi, and O. C. P. Agubosi, "Effect of dietary supplementation of melon (*Citrullus lanatus*) seed oil on the growth performance and antioxidant status of growing rabbits," *J. Multidimensional Res. Rev.*, vol. 2, no. 1, pp. 78–95, 2021.
- [33] M. Hernandez and J. O. Alagbe, "Influence of *Abrus precatorius* crude oil supplementation on growth performance, nutrient digestibility, ruminal fermentation and microbial population of Malabari bucks," *Int. J. Global Sustain. Res.*, vol. 3, no. 7, pp. 527–538, 2025.
- [34] M. Hernandez and J. O. Alagbe, "Influence of *Odontonema strictum* oil on the growth performance and ruminal fermentation of Barbari bucks," *Res. Agric. Vet. Sci.*, vol. 9, no. 2, pp. 1–10, 2025.
- [35] D. S. Muritala, D. Shittu, J. O. Alagbe, O. O. Ojebiyi, T. K. Ojediran, and T. A. Rafiu, "Growth performance and haematological and serum biochemical parameters of broiler chickens given varied concentrations of *Polyalthia longifolia* leaf extract in place of conventional antibiotics," *Anim. Sci. Genet.*, vol. 18, no. 2, pp. 57–71, 2022.
- [36] T. K. Ojediran, I. A. Emiola, V. Durojaye, and J. O. Alagbe, "Analysis of *Kigelia africana* fruit powder antioxidant and phytochemical properties," *Brazilian J. Sci.*, vol. 3, no. 7, pp. 38–49, 2024.
- [37] T. K. Ojediran, I. A. Emiola, V. Durojaye, and A. O. John, "Proximate, vitamin and GC–MS profiling of *Kigelia africana* powder," *Cerrado: Agric. Biol. Res.*, vol. 1, no. 1, pp. 13–20, 2024.
- [38] M. D. Shittu, J. O. Alagbe, O. Alaba, E. O. Okanlawon, F. A. Adelokun, E. O. Emmanuel, and D. O. Adejumo, "Effect of ginger, garlic and negro pepper on gut microbes, gut histomorphometry and pathological assessment of selected organs of broiler chickens," *Assoc. Deans Agric. Niger. Univ.*, vol. 5, pp. 105–121, 2024.
- [39] R. A. Oluwafemi, A. O. Lawal, S. A. Adelowo, and J. O. Alagbe, "Effects of dietary inclusion of ginger (*Zingiber officinale*) and garlic (*Allium sativum*) oil on carcass characteristics and sensory evaluation of broiler chickens," *Texas J. Multidiscip. Stud.*, vol. 2, no. 11, pp. 180–188, 2021.
- [40] J. O. Alagbe, M. D. Shittu, T. Tanimomo, and B. K. Babatunde, "Influence of *Anogeissus leiocarpus* stem bark on fatty acid composition in broiler chicken meat," *Eur. J. Life Safety Stability*, vol. 14, no. 22, pp. 13–22, 2022.
- [41] J. O. Alagbe, "Use of medicinal plants as a panacea to poultry production and food security: A review," *Gospodarka i Innowacje*, vol. 22, pp. 1–12, 2022.
- [42] E. O. Omokore and J. O. Alagbe, "Efficacy of dried *Phyllanthus amarus* leaf meal as an herbal feed additive on growth performance, haematology and serum biochemistry of growing rabbits," *Int. J. Acad. Res. Dev.*, vol. 4, no. 3, pp. 97–104, 2019.
- [43] B. Musa, J. O. Alagbe, M. B. Adegbite, and E. A. Omokore, "Growth performance, caeca microbial population and immune response of broiler chicks fed aqueous extract of *Balanites aegyptiaca* and *Alchornea cordifolia* stem bark mixture," *United J. Res. Technol.*, vol. 2, no. 2, pp. 13–21, 2020.
- [44] A. O. Adewale, J. O. Alagbe, and A. O. Adeoye, "Dietary supplementation of *Rauvolfia vomitoria* root extract as a phytogenic feed additive in growing rabbit diets: Haematology and serum biochemical indices," *Int. J. Orange Technol.*, vol. 3, no. 3, pp. 1–12, 2021.
- [45] O. J. Alagbe, "Novel phytogenics' impact on weaned pigs growth performance, haematology and serum biochemical indicators," *Black Sea J. Agric.*, vol. 7, no. 2, pp. 82–89, 2024.
- [46] M. D. Shittu and J. O. Alagbe, "Phyto-nutritional profiles of broom weed (*Sida acuta*) leaf extract," *Int. J. Integr. Educ.*, vol. 3, no. 11, pp. 119–124, 2020.

- [47] J. O. Alagbe, "Gas chromatography–mass spectroscopy of *Juniperus phoenicea* stem bark extract and its influence on haemato-biochemical values of growing rabbits," *British Sci. Period.*, vol. 1, no. 1, pp. 18–33, 2022.
- [48] J. O. Alagbe, H. Zubairu, O. M. Adedeji, S. Bamigboye, and A. Dora, "Influence of *Juniperus thurifera* root extract on nutrient digestibility and caecal microbial count of growing rabbits," *Web Synergy: Int. Interdiscip. Res. J.*, vol. 1, no. 1, pp. 5–17, 2022.
- [49] National Institute of Standards and Technology (NIST), NIST Chemistry WebBook, Standard Reference Database 69, 2011.
- [50] R. R. Sharma, "Preliminary phytochemical screening of some indigenous medicinal plant leaf extracts in regulation of antidiabetic activity," *Sci. Res. Rep.*, vol. 2, no. 3, pp. 307–310, 2012.
- [51] J.-C. Chou, J.-Y. Yang, P.-C. Chen, and C.-K. Hsu, "Phenolic content and DPPH radical scavenging activity of yam-containing surimi gels influenced by salt and heating," *Asian J. Health Inf. Sci.*, vol. 2, pp. 1–11, 2007.
- [52] A. L. Lima, R. Parial, M. Das, and A. K. Das, "Phytochemical and pharmacological studies of ethanolic extract from the leaf of mangrove plant *Phoenix paludosa* Roxb.," *Malaysian J. Pharm. Sci.*, vol. 8, no. 2, pp. 59–69, 2010.
- [53] A. Manilal et al., "Biological activity of the red algae *Laurencia brandenii*," *Acta Bot. Croat.*, vol. 70, no. 1, pp. 81–90, 2011.
- [54] H. S. Namkeleja, N. T. C. Tarimo, and P. A. Ndakidemi, "Allelopathic effects of *Argemone mexicana* on growth of native plant species," *Am. J. Plant Sci.*, vol. 5, no. 9, pp. 1336–1344, 2014.
- [55] P. Tiwari, B. Kumar, M. Kaur, G. Kaur, and H. Kaur, "Phytochemical screening and extraction: A review," *Int. Pharm. Sci.*, vol. 1, no. 1, pp. 98–106, 2011.
- [56] M. K. Sinha and J. D. Munshi, *Toxicology of Biocidal Plants*. Delhi, India: Mittal Publications, 2010, p. 53.
- [57] H. Zubairu, A. A. Mubarak, and J. O. Alagbe, "Effect of partial replacement of soybean meal with *Sphenocentrum jollyanum* seed meal on growth performance and haematological constituents of broiler chickens," *Int. J. Integr. Res.*, vol. 3, no. 10, pp. 741–752, 2025.
- [58] J. O. Alagbe, "Accessing the bioactive compounds of *Thaumatococcus daniellii* leaf extract by GC–MS," *Int. J. Global Sustain. Res.*, vol. 3, no. 9, pp. 617–626, 2025.
- [59] O. C. P. Agubosi, A. Alexander, and J. O. Alagbe, "Influence of dietary inclusion of sunflower (*Helianthus annuus*) oil on growth performance and oxidative status of broiler chicks," *Cent. Asian J. Med. Nat. Sci.*, vol. 2, no. 7, pp. 187–195, 2022.
- [60] O. C. P. Agubosi, M. B. Soliu, and J. O. Alagbe, "Effect of dietary inclusion levels of *Moringa oleifera* oil on growth performance and nutrient retention of broiler starter chicks," *Cent. Asian J. Theor. Appl. Sci.*, vol. 3, no. 3, pp. 30–39, 2022.