

# Efficacy of Topical Treatment for Acute Rhinosinusitis in Children

Nasretdinova Makhzuna Takhsinovna <sup>1</sup>, Khusanova Dilnoza Abdukhamid qizi <sup>2</sup>,  
Normuradov Nodir Alisherovich <sup>3</sup>

<sup>1,3</sup> Samarkand State Medical University

<sup>2</sup> Termez Branch of Tashkent Medical Academy

**Received:** 2024, 15, Feb

**Accepted:** 2025, 21, Mar

**Published:** 2025, 14, Apr

Copyright © 2025 by author(s) and BioScience Academic Publishing. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).



Open Access

<http://creativecommons.org/licenses/by/4.0/>

**Annotation:** The article is devoted to the problem of acute rhinosinusitis in children. The data of separate researches on estimation of efficiency and expediency of using hypertonic solutions of sea water in complex treatment of rhinosinusitis are presented. It is shown that inclusion of a spray with hypertonic solution of sea water in the treatment scheme allows to improve objective indicators of nasal function and patient's condition in a short period of time, shortens the total duration of the disease, reduces the risk of complications.

**Keywords:** rhinosinusitis, treatment, children's age, hypertonic solutions, seawater.

**Introduction.** Acute rhinosinusitis is one of the most common diseases of the ENT organs in all age groups, ranking 5th among all infections in terms of the number of antibiotics prescribed [1, 2]. The leading places among the causative agents of acute sinusitis in children are occupied by *Streptococcus pneumoniae* - 36%, *Haemophilus influenzae* - 23%, *Moraxella catarrhalis* - 20% [3], and the spectrum of causative agents and their percentage ratio have remained unchanged in recent years. The study of the main causative agents and the characteristics of the clinical course of rhinosinusitis allows us to conclude that in most cases this pathology develops after a previous respiratory viral infection. The average duration of upper respiratory tract diseases during a lifetime is estimated to be 1 to 2 years. In Russia, according to various data, from 10 to 15 million people fall ill with rhinosinusitis annually. In the first stage of the disease, the virus adheres to the mucous membrane of the upper respiratory tract. The outcome of this contact (the development of an inflammatory reaction or the elimination of the pathogen) depends on the state of the epithelium of the upper respiratory tract. That is why the issues of rational treatment and prevention of upper respiratory tract respiratory infections (and their complications) are closely related to the normalization of the functioning of the ciliated epithelium, which is impaired in inflammation of any etiology [4–8]. Various groups of drugs can be used to

normalize mucociliary clearance and eliminate the inflammatory process. Taking into account the etiopathogenesis, antiviral or antibacterial (etiotropic) drugs occupy a leading place in this list. However, one should not forget about the sensitivity (growth of resistance) of causative pathogens to this group of drugs.

Widely used in topical therapy of rhinosinusitis

- ✓ irrigation-elimination means for therapy;
- ✓ topical decongestants;
- ✓ steroid anti-inflammatory drugs;
- ✓ topical mucolytic drugs.

In the treatment and prevention of diseases of the upper respiratory tract irrigation-elimination. The experience of using saline nasal irrigation goes back thousands of years and is fully justified. Since the time of Hippocrates, nasal irrigation with saline solutions has been considered an important stage in the local treatment of nasal diseases. The results of numerous modern studies indicate the positive effect of moistening the mucous membrane on mucociliary activity, as well as the safety of the procedure, which has no documented serious side effects, which allows it to be used in the treatment of rhinosinusitis in children [9, 10]. Nasal irrigation with saline solutions helps to reduce the impact of various microorganisms and harmful particles in the inhaled air on the nasal mucosa, making it difficult to resorb toxins and allergens. Isotonic, hypo- and hypertonic solutions of sodium chloride are used in daily practice for nasal irrigation. These solutions dilute mucus and regulate its production by goblet cells, soften the drying secretion, improve blood circulation in the mucous membrane, increase its sensitivity and reactivity. The salt concentration in the isotonic solution is close to the salt concentration in the body's fluid environments, therefore it has a mild therapeutic effect on the nasal mucosa, contributing to the normalization of the hydro-ionic balance, the violation of which is accompanied by swelling of the nasal and sinus mucosa. Hypertonic saline has a higher salt concentration, which leads to an accelerated movement of water towards a higher concentration of ions and has a pronounced anti-edema and stimulating effect. Acting as a stimulant on the nasal mucosa, hypertonic saline stimulates nociceptive nerves and glandular secretion without changing vascular permeability [11].

Hypertonic saline has been shown to restore ciliary epithelial function more effectively in patients with chronic sinusitis, while isotonic saline has been shown to improve mucociliary transport in patients with allergic rhinitis and acute sinusitis [12]. This also applies to seawater, which is similar in composition to chloride-sodium mineral waters. The beneficial effects of seawater, which is predominantly sodium chloride (more than 70% of all salts) on the nasal mucosa, are due to the unique properties of its physicochemical composition. Under the influence of salt, the production of nasal mucus by goblet cells is normalized, and nasal secretions are liquefied. Microelements in seawater (mainly Ca, Fe, K, Mg, Cu) lead to the normalization of the function of the ciliated epithelium, an increase in the resistance of the nasal mucosa to infection, and an improvement in the subjective sensations of patients. When comparing the effects of hypertonic solutions of Dead Sea salts and hypertonic saline solutions of sodium chloride on the quality of life of patients with chronic rhinosinusitis, it was shown that both groups of patients improved, but the quality of life indicators were better in the use of seawater [13, 14].

**Research purpose** was to evaluate the effectiveness of hypertonic solutions in the treatment of acute rhinosinusitis in children.

**Materials and methods of the study.** 100 children aged 2-15 years, who were treated in outpatient or inpatient conditions at the Department of Otorhinolaryngology, Multidisciplinary Clinic of Samarqand State Medical University, with severe nasal congestion, abundant mucous

or mucopurulent discharge from the nose, itching and maceration in the nasal passages, decreased sense of smell, and periodic dryness in the nasopharynx, were examined. In addition to the standard otorhinolaryngological clinical examination, the examination regimen of patients included X-ray of the nasal cavity, endoscopy, and previous active rhinomanometry.

One of the preparations based on seawater is the hypertonic AQUALOR forte ("PharmaMed", Canada). This spray is a natural sterile hypertonic seawater (salt content - 21 g/l) intended for washing the nasal cavity, which contains all the active substances and trace elements of seawater (Mg, Zn, Se, Cu, Fe, I, Ca, Na, etc.) necessary to maintain the normal physiological state of the nasal mucosa. The spray is equipped with a nozzle that provides a powerful shower spray. The efficacy and tolerability of this drug in the complex treatment of acute uncomplicated rhinosinusitis were evaluated in an open comparative randomized study involving children aged 2 to 15 years who received a course of treatment for acute uncomplicated rhinosinusitis in an outpatient or inpatient setting. The diagnosis was made on the basis of complaints, physical examination and special examination methods (nasal radiography, endoscopy).

In addition to the subjective assessment of the severity of acute rhinosinusitis symptoms, objective monitoring was performed, which included the following:

- endoscopic examination of the nasopharynx (presence of hyperemia and edema in the mucous membrane of the nasopharynx, the nature of the discharge);
- previous active rhinomanometer (allows to evaluate the ventilatory function of the nose).

The essence of active rhinomanometry is to measure nasal resistance (based on quantitative measurement of nasal airflow and pressure). Nasal resistance is measured in pascals (PA), volumetric flow in cubic centimeters per second (cm<sup>3</sup>/s). The results are given in the International System of Units (SI): Pa/cm<sup>3</sup> per second. All results were entered into the patient's individual card. 100 children were observed, divided into 2 groups. The distribution of subjects into groups was carried out according to a table of random numbers (created using the standard function of the Statistica 6 statistical software package).

Group 1 (main) - consisted of 50 patients who, in addition to the standard therapy of the disease (topical decongestants 3 times a day and antibacterial drugs taken orally according to the scheme), received daily (at least 4-6 times a day) nasal irrigation with AQUALOR forte hypertensive spray for 7-10 days. Group 2 (comparison) - consisted of 50 patients who received only standard treatment (antibiotic and topical decongestant).

exacerbation of chronic sinusitis or complicated acute sinusitis were excluded from the study. The following criteria were used to assess the effectiveness of treatment:

- ✓ recovery — complete disappearance of clinical symptoms of the disease;
- ✓ improvement — a significant reduction in the expression of clinical symptoms of the disease;
- ✓ lack of effect - lack of positive dynamics, exacerbation of disease symptoms.

**Inspection results.** Before starting treatment, all patients had similar complaints: severe nasal congestion, copious mucous or mucopurulent nasal discharge, itching and maceration in the nasal vestibule, decreased sense of smell, and periodic dryness in the nasopharynx. Previous rhinoscopy revealed hyperemia and edema of the nasal mucosa, severe narrowing of the common nasal passages, and copious mucous or mucopurulent discharge. Subjective and objective assessments of acute uncomplicated rhinosinusitis in the observed patients before, during, and after treatment are presented in Table 1. In the main group of children who received hypertensive spray in addition to standard therapy (antibiotics and topical decongestants), a decrease in the severity of clinical symptoms was noted on the 3rd day after starting treatment. In 54% of patients (according to parents), a significant decrease in dryness of the nasal and nasopharyngeal mucosa was observed. On the 3rd-4th day of treatment, subjective improvement of the condition was noted in 76% of patients. The rhinoscopic appearance of patients in this

group was manifested by a significant decrease or disappearance of edema or hyperemia of the nasal mucosa on the 5th-7th day of treatment, the absence of mucous discharge from the nasal cavity, and the restoration of the sense of smell. Also, the frequency of use of topical decongestants in the main group decreased: already on the 3rd day of treatment, 100% of children used these drugs only before bedtime, which was enough to ensure free nasal breathing throughout the day.

**Table 1. Expression of symptoms of acute uncomplicated rhinosinusitis in children before and after treatment on a 10-point visual analogue scale**

symptom	1 -day		2-day		3-day	
	main	control	main	control	main	control
Increased body temperature	4	4	3	4	1	2
Weakness	6	6	4	5	1	2
Intoxication	5	5	4	4	1	2
It is not separated from the nose	6	6	3	4	0	1
Nasal breathing disorder	6	6	4	6	2	3
membrane g hyperemia	6	6	3	5	1	3
Flow along the posterior wall of the larynx	5	5	2	3	0	1
The differences are huge	6	6	2	3	0	1

#### Literature used:

1. Козлов В. С., Шиленкова В. В., Шиленков А. А. Синуситы: современный взгляд на проблему //Consilium medicum. – 2003. – Т. 5. – №. 4. – С. 212-218.
2. Насретдинова М. Т., Хайитов А. А. Усовершенствованный подход при лечении кист верхнечелюстной пазухи.
3. Насретдинова М. Т., Махмудова Н. Р., Хайитов А. А. СОВЕРШЕНСТВОВАНИЕ ДИАГНОСТИКИ У ДЕТЕЙ С ХРОНИЧЕСКИМ ГНОЙНЫМ РИНОСИНУСИТОМ //The 5th International scientific and practical conference “Modern scientific research: achievements, innovations and development prospects”(October 24-26, 2021) MDPC Publishing, Berlin, Germany. 2021. 686 p. – 2021. – С. 122.
4. Насретдинова М., Хайитов А. Юқори жағ бўшлиғининг кисталарида тежамкор эндоназал олиб ташлаш ҳақида //Журнал вестник врача. – 2021. – Т. 1. – №. 1 (98). – С. 94-96.
5. Насретдинова М. Т., Хайитов А. А., Нормурадов Н. А. Состояние микробиоциноза у пациентов скистозным поражением верхнечелюстных синусов //Оториноларингология. Восточная Европа. – 2021. – Т. 11. – №. 2. – С. 169-174. Насретдинова М. и др. О ЩАДЯЩЕМ ЭНДОНАЗАЛЬНОМ УДАЛЕНИИ КИСТ ВЕРХНЕЧЕЛЮСТНЫХ ПАЗУХ //Журнал стоматологии и краниофациальных исследований. – 2020. – Т. 1. – №. 3. – С. 24-26.
6. Насретдинова М., Хайитов А. Наша тактика при хирургическом лечении кистозных поражений гайморовых пазух //Журнал вестник врача. – 2020. – Т. 1. – №. 2. – С. 72-75.

7. Насретдинова, М., & Хайитов А. (2022). О щадящем эндоназальном удалении кист верхнечелюстных пазух. Журнал вестник врача, 1(1), 94–96.
8. Шиленкова В. В. Острые и рецидивирующие синуситы у детей (диагностика и лечение) //Автореф. дисс. д. м. н., М. – 2008.
9. Nasretdinova Makhzuna Takhsinovna, Khayitov Alisher Adkhamovich, Buriyev Shahzod Saidulla ugli, & Lapasov Mirsaid Hassan ugli. (2024). THE DIAGNOSIS AND TREATMENT CYSTS OF MAXILLARY SINUS. Web of Scientist: International Scientific Research Journal, 5(2), 62–65
10. Taxsinovna, N. M. ., Adxamovich, X. A. ., Ibragimovna, R. E. ., & Sherbek ugli, R. S. . (2024). A Modern Approach to the Diagnosis and Treatment of Paranasal Sinus Cysts. Research Journal of Trauma and Disability Studies, 3(2), 149–152.
11. Takhsinovna N. M., Musinovna R. K. CHARACTERISTICS OF COMPLAINTS FROM WORKERS IN NOISE PROFESSIONS, HAVING HEARING IMPAIRMENTS //Health Horizon: Congress on Public Health and Biomedical Sciences. – 2025. – Т. 1. – №. 1. – С. 28-30.
12. Taxsinovna N. M. et al. VESTIBULAR VASCULAR REACTIONS IN ASSESSMENT OF VESTIBULAR DYSFUNCTION IN PATIENTS WITH CRANIOCEREBRAL TRAUMA //INNOVATIVE ACHIEVEMENTS IN SCIENCE 2024. – 2024. – Т. 3. – №. 29. – С. 104-111.